



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A CATALOGUE OF THE FISHES OF BEAN BLOSSOM CREEK, MONROE
COUNTY, INDIANA.

BY CARL H. EIGENMANN AND MORTON W. FORDICE.

Bean Blossom is a small creek crossing Monroe County about six miles north of Bloomington. It is a tributary of White River and empties into it near Gosport, Indiana. It is a rather sluggish stream with gravelly bottom, and considerable grass and water-weeds. The specimens were collected by parties of students of the Indiana University, visiting the creek at different times. Thirty-two of the species were also obtained by us on the 12th of September, 1885.

1. *Ammocætes branchialis* L. Abundant in spring.
2. *Noturus gyrinus* Mitchill.
3. *Noturus miurus* Jordan. Very abundant.
4. *Noturus flavus* Rafinesque. Very common. The last two species are very abundant in the weeds of shallow water; only two specimens of *gyrinus* were obtained.
5. *Noturus exilis* Nelson. One specimen.
6. *Leptops olivaris* Rafinesque.
7. *Amiurus melas* Rafinesque.
8. *Amiurus natalis* Le Sueur.
9. *Catostomus teres* Mitchill. Very abundant.
10. *Catostomus nigricans* Le Sueur. Abundant.
11. *Mnytrema melanops* Rafinesque. Abundant.
12. *Moxostoma macrolepidotum* Le Sueur.
13. *Campostoma anomalum* Rafinesque.
14. *Chrosomus erythrogaster* Rafinesque.
15. *Hybognathus nuchalis* Agassiz.
16. *Notropis whipplei* Girard.
17. *Notropis megalops* Rafinesque.
18. *Notropis ardens lythrurus* Jordan.
19. *Rhinichthys atronasus* Mitchill.
20. *Hybopsis biguttatus* Kirtland.
21. *Hybopsis amblops* Rafinesque.
22. *Semotilus atromaculatus* Mitchill.
23. *Esox vermiculatus* Le Sueur.
24. *Labidesthes sicculus* Cope. One specimen.
25. *Aphredoderus sayanus* Gilliams. One specimen.
26. *Pomoxys annularis* Rafinesque. Very abundant.
27. *Pomoxys sparoides* Lacépède.
28. *Ambloplites rupestris* Rafinesque.
29. *Lepomis cyanellus* Rafinesque.

30. *Lepomis megalotis* Rafinesque. Abundant.
31. *Micropterus dolomieu* Lacépède. Abundant.
32. *Boleosoma olmstedii maculatum* Agassiz. Abundant.
33. *Diplexion blennioides* Rafinesque. Abundant.
34. *Percina caprodes* Rafinesque. Abundant.
35. *Hadropterus phoxocephalus* Nelson.
36. *Hadropterus scierus* Swain. Abundant. Several specimens were taken five inches in length. In these large specimens the serration of the preopercle is obsolete.
37. *Hadropterus aspro* Cope and Jordan.
38. *Etheostoma flabellare* Rafinesque. Abundant.
39. *Etheostoma caeruleum* Storer.
40. *Uranidea richardsoni* Agassiz.